

# OXI PA

**NRCS Registered -**

Oxi PA is a 15-17% active peracetic acid based liquid disinfectant. Suitable for use in CIP systems, dairy tanks as well as stainless steel tanks. Oxi PA is predominately used in the Health Care Industry for disinfecting and destaining in wash tunnels. Product also neutralizes offensive odours in industrial laundries. Product pH - undiluted is between 1.00 - 3.00.

## Dilutions

Use 1:10 - 1:100 dilution with water, depending on application. Allow a 5 minute contact time. Use at 1:100 for bacteria, and 1:20 for Moulds and Spores. Ensure proper PPE is worn at all times. Do not allow contact with Skin or Eyes. Avoid breathing in fumes. Use diluted product within 24 hours.

## Microbial Efficacy

Bactericidal – Log 7 – 1:100 with a 5 min contact time - *Pseudomonas aeruginosa*.

Bactericidal – Log 6 @ 1:20 with a 5 min contact time - *Listeria monocytogenes*

Fungicidal - Log 4 – @ 1:20 with a 15 min contact time - *Aspergillus brasiliensis*

Sporicidal - Log 4 - @ 1:20 with 6 hours contact time - *Bacillus subtilis spores*

Sporicidal - Log 4 - @ 1:20 with 4 hours contact time - *Bacillus subtilis spores*

## GHS Identification:

Danger

**UN 3109 – organic Peroxides, Type F, Liquid**



## PPE

Gloves, Goggles, Personal Protective Clothing, Apron, Gumboots



First Aid

SKIN	EYES	IINHALATION	ORAL
Rinse off with plenty of water for 15 minutes.	Rinse cautiously with water for several minutes, remove contact lenses if safe and easy to do so, continue rinsing and get medical attention	If INHALED remove person to fresh air and keep comfortable for breathing.	Do NOT induce vomiting
Causes severe burns to skin. Consult a Doctor is pain persists.			Consult a Doctor if you feel unwell
			Consult a Doctor if feeling unwell

Available In  
25Lt